arXiv:1303.6019v1 [math.DG] 25 Mar 2013

Perelman's entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials

Songzi Li, Xiang-Dong Li*

March 26, 2013

Abstract. In this paper, we first give a new proof of the W-entropy formula for the heat equation associated with the Witten Laplacian on compact Riemannian manifolds with a fixed metric and a fixed potential. This gives a natural geometric interpretation of a quantity appeared in the W-entropy formula. Then we extend the W-entropy formula to the heat equation of the Witten Laplacian on compact Riemannian manifolds with time dependent metrics and potentials. As an application, we prove that the optimal logarithmic Sobolev constant associated with the Witten Laplacian on compact manifolds equipped with the m-dimensional Perelman's super Ricci flow is decreasing in time. Finally, we derive the W-entropy formula for the backward heat equation associated with the Witten Laplacian on compact Riemannian manifolds equipped with Lott's modified Ricci flow.

1 Introduction

Let M be a complete Riemannian manifold with a fixed Riemnnian metric g and a fixed potential $\phi \in C^2(M)$. Let $d\mu = e^{-\phi}dv$, where dv is the Riemannian volume measure on (M, g). The Witten Laplacian, called also the weighted Laplacian,

$$L = \Delta - \nabla\phi \cdot \nabla$$

is a self-adjoint and non-positive definite operator on $L^2(M, \mu)$. By Itô's calculus, one can construct the symmetric diffusion process X_t associated to the Witten Laplacian by solving the SDE

$$dX_t = \sqrt{2}dW_t - \nabla\phi(X_t)dt,$$

where W_t is the Brownian motion on M. Moreover, it is well known that the transition probability density function of the diffusion process X_t is exactly the fundamental solution to the heat equation of L, i.e., the heat kernel of the Witten Laplacian L. In view of this, it is a fundamental problem to study the heat equation and the heat kernel of the Witten Laplacian on manifolds.

In recent years, important progress has been obtained in the study of the heat equation associated with the Witten Laplacian by using new ideas and new methods from geometric analysis, PDEs, and probability theory. In particular, F. Otto [26] introduced an infinite dimensional Riemannian structure on the Wasserstein space of probability measures on $\mathbb{R}^n$ and proved that the heat equation

$$\partial_t u = Lu \tag{1}$$

---

*Research supported by NSFC No. 10971032, Key Laboratory RCSDS, CAS, No. 2008DP173182, and a Hundred Talents Project of AMSS, CAS.

can be realized as the reverse gradient flow of the Boltzmann-Shannon entropy[1]

$$H(u) = -\int_M u \log u d\mu.$$

See also [27, 32, 33, 34, 35] for the extension of Otto's work to Riemannian manifolds.

The Witten Laplacian is a natural extension of the standard Laplace Beltrami operator and has a close connection with differential geometry, probability theory, quantum field theory and statistical mechanics. In view of this, it is natural to raise the question whether one can extend the results which hold for the standard Laplace Beltrami operator to the Witten Laplacian on manifolds. The main tool which leads such an extension possible is the so-called Bakry-Emery Ricci curvature associated to L introduced in [2], i.e.,

$$Ric(L) = Ric + \nabla^2 \phi,$$

which plays the same rôle as the Ricci curvature for the standard Laplace Beltrami operator. We refer the reader to [4, 3, 17] for the Li-Yau Harnack estimates and the heat kernel estimates to the heat equation (1), and to [17] for the extension of S.-T. Yau's Strong Liouville theorem for the positive L-harmonic functions and the L^1-uniqueness of the heat equation (1) on complete Riemannian manifolds. See also [1, 5, 12, 13, 27, 33, 34, 35, 36] for further works on the study of the Witten Laplacian and Bakry-Emery Ricci curvature on manifolds with weighted measures.

The Bakry-Emery Ricci curvature has been essentially used in Perelman's work [30] on the entropy formula for Ricci flow, where Perelman first introduced the $\mathcal{F}$-energy functional on $\mathcal{M} = \{\text{all Riemannian metrics } g \text{ on } M\} \times C^\infty(M)$ as follows

$$\mathcal{F}(g, f) = \int_M (R + |\nabla f|^2) e^{-f} dv,$$

where $f \in C^\infty(M)$, R denotes the scalar curvature on (M, g), and dv denotes the volume measure. Under the constraint condition which requires that

$$d\mu = e^{-f} dv$$

is a fixed weighted measure on (M, g), Perelman [30] proved that the gradient flow of the $\mathcal{F}$-energy functional with respect to the standard L^2-Riemannian metric on $\mathcal{M}$ is given by the following modified Ricci flow, which we call Perelman's Ricci flow in the sequel,

$$\partial_t g = -2(Ric + \nabla^2 f),$$

and f satisfies the so-called conjugate heat equation

$$\partial_t f = -\Delta f - R.$$

Moreover, Perelman [30] introduced the W-entropy functional and proved its monotonicity for the Ricci flow on compact manifolds. This result plays an important rôle in the proof of the no local collapsing theorem and in the final resolution of the Poincaré conjecture and geometrization conjecture (see e.g. [7, 25, 15]). Since then, many people have derived the W-entropy formula for various geometric evolution equations and used it to study further analysis and geometric properties of manifolds. See e.g. [10, 8, 15, 28, 29, 11, 23, 24].

In [18], see also [19, 20], inspired by Perelman's work on the W-entropy formula for Ricci flow, the second author proved the W-entropy formula for the fundamental solution of the Witten-Laplacian on complete Riemannian manifolds with bounded geometry condition, which extends a previous result due to Ni [28, 29], who proved an analogue of Perelman's W-entropy formula for the heat equation $\partial_t u = \Delta u$ on complete Riemannian manifolds with a fixed metric. To be more precise, let

---

[1] Equivalently, the heat equation (1) is the gradient flow of $Ent(u) = -H(u)$ on the Wasserstein space $\mathscr{P}_2(\mathbb{R}^n)$ equipped with Otto's infinite dimensional Riemannian metric.

$$Ric_{m,n}(L) = Ric + \nabla^2\phi - \frac{\nabla\phi \otimes \nabla\phi}{m-n}$$

be the m-dimensional Bakry-Emery Ricci curvature associated with the Witten Laplacian L. Then we have the following result.

Theorem 1.1 *([18, 19, 20]) Let (M, g) be a complete Riemannian manifold with bounded geometry condition*[2]*, and $\phi \in C^4(M)$ with $\nabla\phi \in C_b^3(M)$. Let $m > n$, and $u = \frac{e^{-f}}{(4\pi t)^{m/2}}$ be the fundamental associated with the Witten Laplacian, i.e., the heat kernel to the heat equation $\partial_t u = Lu$. Let*

$$H_m(u, t) = -\int_M u \log u d\mu - \frac{m}{2}(1 + \log(4\pi t)),$$

and define

$$W_m(u, t) := \frac{d}{dt}(tH_m(u)). \tag{2}$$

Then

$$W_m(u, t) = \int_M \left[t|\nabla \log u|^2 + f - m\right] \frac{e^{-f}}{(4\pi t)^{m/2}} d\mu,$$

and

$$\begin{aligned} \frac{dW_m(u,t)}{dt} &= -2\int_M t\left(\left|\nabla^2 f - \frac{g}{2t}\right|^2 + Ric_{m,n}(L)(\nabla f, \nabla f)\right) u d\mu \\ &\quad -\frac{2}{m-n}\int_M t\left(\nabla\phi \cdot \nabla f + \frac{m-n}{2t}\right)^2 u d\mu. \end{aligned}$$

In particular, if (M, g, ϕ) satisfies the bounded geometry condition and $Ric_{m,n}(L) \geq 0$, then the W-entropy is decreasing in time t, i.e.,

$$\frac{dW_m(u,t)}{dt} \leq 0, \qquad \forall t \geq 0.$$

The purpose of this paper is to extend the W-entropy formula in Theorem 1.1 to the heat equation (1) associated with the time dependent Witten Laplacian on compact Riemannian manifolds equipped with time dependent metrics and potentials. In view of Perelman's work using the W-entropy formula for the Ricci flow to remove "the major stumbling block in Hamilton's approach to geometrization" ([30]), it might be possible that the W-entropy formula for the time dependent Witten Laplacian can bring some new insights to the study of geometric analysis on Riemannian manifolds with time dependent metrics and potentials.

We now in a position to state the main result of this paper as follows.

Theorem 1.2 *Let $(M, g(t), t \in [0, T])$ be a family of compact Riemannian manifolds with potential functions $f(t) \in C^\infty(M)$, $t \in [0, T]$. Suppose that $g(t)$ and $f(t))$ satisfies the conjugate equation*

$$\frac{\partial f}{\partial t} = \frac{1}{2}\mathrm{Tr}\left(\frac{\partial g}{\partial t}\right).$$

---

[2] Here we say that (M, g) satisfies the bounded geometry condition if the Riemannian curvature tensor Riem and its covariant derivatives ∇^kRiem are uniformly bounded on M, $k = 1, 2$.

Let

$$L = \Delta_{g(t)} - \nabla_{g(t)} f(t) \cdot \nabla_{g(t)}$$

be the time dependent Witten Laplacian on $(M, g(t), f(t))$*. Let* u *be a positive solution of the heat equation*

$$\partial_t u = Lu$$

with initial date $u(0)$ *satisfying* $\int_M u(0)d\mu(0) = 1$*. Let*

$$H_m(u,t) = -\int_M u \log u d\mu - \frac{m}{2}(1 + \log(4\pi t)),$$

and define

$$W_m(u,t) := \frac{d}{dt}(tH_m(u)).$$

Then

$$W_m(u,t) = \int_M \left[t|\nabla \log u|^2 - \log u - m\right] ud\mu,$$

and

$$\begin{aligned}\frac{d}{dt}W_m(u,t) \quad &= \quad -2t\int_M \left|\nabla^2 \log u + \frac{g}{2t}\right|^2 ud\mu - \frac{2t}{m-n}\int_M \left(\nabla f \cdot \nabla \log u - \frac{m-n}{2t}\right)^2 ud\mu \\ &\quad -2\int_M t\left(\frac{1}{2}\frac{\partial g}{\partial t} + Ric_{m,n}(L)\right)(\nabla \log u, \nabla \log u)ud\mu.\end{aligned}$$

In particular, if $\{g(t), f(t), t \in (0,T]\}$ *satisfies the* m*-dimensional Perelman's super Ricci flow and the conjugate equation*

$$\frac{1}{2}\frac{\partial g}{\partial t} + Ric_{m,n}(L) \geq 0, \tag{3}$$

$$\frac{\partial f}{\partial t} = \frac{1}{2}\mathrm{Tr}\left(\frac{\partial g}{\partial t}\right), \tag{4}$$

then $W_m(u,t)$ *is decreasing in* $t \in (0,T]$*, i.e.,*

$$\frac{dW_m(u,t)}{dt} \leq 0, \quad \forall t \in (0,T].$$

As an application of the W-entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials, we prove that the optimal logarithmic Sobolev constant associated with the Witten Laplacian on compact manifolds equipped with the m-dimensional Perelman's super Ricci flow is decreasing in time. More precisely, we have

Theorem 1.3 *Let* $(M, g(t), f(t), t \in [0,T])$ *be as in Theorem 1.2. Then there exists a positive and smooth function* $u = e^{-\frac{v}{2}}$ *such that* v *achieves the optimal logarithmic Sobolev constant* $\mu(t)$ *defined by*

$$\mu(t) := \inf\left\{W_m(u,t) : \int_M \frac{e^{-v}}{(4\pi t)^{m/2}}d\mu = 1\right\}.$$

Indeed, $u = e^{-\frac{v}{2}}$ *is a solution to the nonliniear PDE*

$$-4tLu - 2u\log u - mu = \mu(t)u.$$

Moreover, if $\{g(t), f(t), t \in [0,T]\}$ *satisfies the* m*-dimensional Perelman's super Ricci flow* (3) *and the conjugate equation* (4)*, then* $\mu(t)$ *is decreasing in* t *on* $[0,T]$*.*

Remark 1.4 In [16], Li and Yau proved the Li-Yau Harnack inequality

$$\Delta \log u + \frac{n}{2t} \geq 0$$

for any positive solution to the heat equation $\partial_t u = \Delta u$ on complete Riemannian manifolds with non-negative Ricci curvature. In general, it was proved in [17, 18] that for any positive solution to the heat equation $\partial_t u = Lu$, the Li-Yau Harnack inequality

$$L \log u + \frac{m}{2t} \geq 0$$

holds on complete Riemannian manifolds with $Ric_{m,n}(L) \geq 0$. See also [4, 3]. By [18], we have the following entropy formula

$$\frac{d}{dt} H_m(u,t) = -\int_M \left(L \log u + \frac{m}{2t} \right) u d\mu. \tag{5}$$

Therefore, $H_m(u,t)$ is decreasing in time t provided that $Ric_{m,n}(L) \geq 0$, i.e.,

$$\frac{d}{dt} H_m(u,t) \leq 0.$$

Remark 1.5 In [30], Perelman gave an interpretation of the W-entropy using the Boltzmann entropy formula in statistical mechanics. In [18, 19], the second author gave a probabilistic interpretation of the W-entropy for the Ricci flow, the heat equation of the Witten Laplacian and for the Fokker-Planck heat equation. Note that, as noticed in [18, 19, 20],

$$H_m(u,t) = H(u,t) - H(\gamma, t)$$

where $H(u,t)$ is the Boltzmann-Shannon entropy associated with the heat equation to the Witten Laplacian on $(M, g(t), f(t))$, and $H(\gamma, t)$ is the Boltzmann-Shannon entropy of the Gaussian heat kernel $\gamma(x,t)$ on $\mathbb{R}^m$ if $m \in \mathbb{N}$ with $m \geq n$, i.e.,

$$\gamma(x,t) = \frac{1}{(4\pi t)^{m/2}} e^{-\frac{|x|^2}{4t}}, \quad x \in \mathbb{R}^m, t > 0.$$

In view of the definition formula (20), the W-entropy $W_m(u,t)$ can be regarded as the by-product of the Boltzmann-Shannon entropy. This gives a probabilistic interpretation of the W-entropy $W_m(u,t)$.

On the other hand, similarly to Perelman [30], we can also give a heuristical interpretation of the W-entropy using the Boltzmann entropy formula in statistical mechanics: Suppose that there exists a canonical ensemble with a "density of state measure" $g(E)dE$ such that the partition function $Z_\beta = \int_{\mathbb{R}^+} e^{-\beta E} g(E) dE$ is given by

$$\log Z_\beta = H_m(u,t), \tag{6}$$

where $t = \beta^{-1}$. Here, as in [30], we do not discuss the issue whether such a "density of state measure" exists or not. Then, formally applying the Boltzmann entropy formula in statistical mechanics, the thermodynamical entropy of this canonical ensemble is given by

$$S = \log Z_\beta - \beta \frac{\partial}{\partial \beta} \log Z_\beta.$$

Using the fact $\frac{\partial}{\partial \beta} = \frac{\partial}{\partial t} \frac{\partial t}{\partial \beta} = -\frac{1}{\beta^2} \frac{\partial}{\partial t} = -t^2 \frac{\partial}{\partial t}$, we have

$$S = H_m(u,t) + t \frac{\partial}{\partial t} H_m(u,t).$$

Thus

$$S = W_m(u,t).$$

Moreover, as noticed in [30], formally using the formula in statistical mechanics

$$\frac{dS}{d\beta} = -\beta \frac{\partial^2}{\partial \beta^2} \log Z_\beta,$$

we can reprove the W-entropy formula in Theorem 1.2.

Remark 1.6 Developing the approach used in [18, 19, 20], it would be possible to further extend the W-entropy formula in Theorem 1.2 to the fundamental solution of the heat equation associated with the Witten Laplacian on complete Riemannian manifolds with time dependent metrics and potentials satisfying the bounded geometry condition. Technically, this requires to prove some Hamilton type gradient estimates (see [14]) for the logarithmic of the heat kernel of the Witten Laplacian on complete Riemannian manifolds with time dependent metrics and potentials satisfying the uniformly bounded geometry condition[3]. We will study this problem in a forthcoming paper. If this can be verified, we can derive that, if $\{g(t), f(t), t \in (0,T]\}$ is a family of metrics and potentials satisfying (3) and (4) on a complete Riemanian manifold M with uniformly bounded geometry condition, then

$$\frac{d}{dt} W_m(u,t) = 0, \qquad \text{for some } \ t = \tau \in (0,T],$$

if and only if

$$\begin{aligned} \nabla^2 \log u &= -\frac{g}{2\tau}, \\ \nabla f \cdot \nabla \log u &= \frac{m-n}{2\tau}. \end{aligned}$$

By the same argument as used in [18, 19, 20], we can further prove the following rigidity theorem: Let $\{g(t), f(t), t \in (0,T]\}$ be a family of metrics and potentials satisfying (3) and (4) on a complete Riemanian manifold M with uniformly bounded geometry condition. Let u be the fundamental solution to the heat equation $\partial_t u = Lu$. Then,

$$\frac{d}{dt} W_m(u,t) = 0, \qquad \text{for some } \ t = \tau \in (0,T],$$

if and only if $(M, g(t))$ is isometric to $\mathbb{R}^n$, $f(t)$ is identically equal to a constant, $m = n$, and

$$u(x,t) = \frac{e^{-\frac{\|x\|^2}{4t}}}{(4\pi t)^{n/2}}, \quad \forall \, x \in M = \mathbb{R}^n, \quad t \in (0,T].$$

The rest of this paper is organized as follows. In Section 2, we first give a new proof[4] to Theorem 1.1. In Section 3, we prove the dissipation formula of the Boltzmann-Shannon entropy for the heat equation of the Witten Laplacian on compact manifolds with time dependent metrics and potentials. In Section 4, we prove Theorem 1.2 and Theorem 1.3. In Section 5, we use Perelman's W-entropy formula for Ricci flow to derive the W-entropy formula for the backward heat equation of the Witten Laplacian on compact Riemannian manifolds equipped with a modified Ricci flow introduced by Lott [22].

---

[3] Here we say that $(M, g(t), f(t), t \in [0,T])$ satisfies the uniformly bounded geometry condition if there exists some $N \in \mathbb{N}$ such that for all $\varepsilon \in (0,T)$, the k-th order covariant derivatives of the Riemannian curvature tensor Riem(g(t)), i.e., ∇^kRiem(g(t)), as well as the k-th order covariant derivatives of $f(t)$, i.e., $\nabla^k f(t)$ are uniformly bounded on $[\varepsilon, T] \times M$, $k = 0, ..., N$.

[4] One of the advantages of our new proof is that it gives a natural geometric interpretation of the third term appeared in the W-entropy formula (3). See Remark 2.2.

2 A new proof of Theorem 1.1

To give a new proof of Theorem 1.1, we first recall some elementary geometric formulas on warped product metrics.

Let $m \in \mathbb{N}$, $m \geq n$. Let $\widetilde{M} = M \times N$, where (N, g_N) is a compact Riemannian manifold with dimension $q = m - n$. Let $\phi \in C^2(M)$. We consider the following warped product metric on $\widetilde{M}$:

$$\widetilde{g} = g_M \bigoplus e^{-\frac{2\phi}{q}} g_N. \tag{7}$$

Let ν_N be the volume measure on N. Then the volume measure on $(\widetilde{M}, \widetilde{g})$ is given by

$$dvol_{\widetilde{M}} = e^{-\phi} dvol_M \otimes d\nu_N.$$

Denote

$$d\mu = e^{-\phi} dvol_M.$$

Then

$$dvol_{\widetilde{M}} = d\mu \otimes d\nu_N.$$

Without loss of generality, we may assume that

$$\nu_N(N) = 1.$$

Let $\widetilde{\Gamma}$ be the Christoffel symbol on $(\widetilde{M}, \widetilde{g})$. By direct calculation, we can verify that

$$\widetilde{\Gamma}_{ij}^k = \Gamma_{ij}^k, \quad \widetilde{\Gamma}_{\alpha\beta}^k = q^{-1} g^{kl} \partial_l \phi g_{\alpha\beta}, \quad \widetilde{\Gamma}_{\alpha\beta}^\gamma = \Gamma_{\alpha\beta}^\gamma,$$

and

$$\widetilde{\Gamma}_{ij}^\alpha = 0, \quad \widetilde{\Gamma}_{i\alpha}^k = 0, \quad \widetilde{\Gamma}_{i\alpha}^\beta = 0.$$

Let $\widetilde{\nabla}$ be the Levi-Civita connection on $(\widetilde{M}, \widetilde{g})$. For any $f \in C^2(M)$, using the formula

$$\begin{aligned}
\widetilde{\nabla}_{ij}^2 f &= \partial_i \partial_j f - \widetilde{\Gamma}_{ij}^k \partial_k f, \\
\widetilde{\nabla}_{i\alpha}^2 f &= \partial_i \partial_\alpha f - \widetilde{\Gamma}_{i\alpha}^k \partial_k f, \\
\widetilde{\nabla}_{\alpha\beta}^2 f &= \partial_\alpha \partial_\beta f - \widetilde{\Gamma}_{\alpha\beta}^k \partial_k f,
\end{aligned}$$

we have

$$\widetilde{\nabla}_{ij}^2 f = \nabla_{ij}^2 f, \tag{8}$$

$$\widetilde{\nabla}_{\alpha\beta}^2 f = -q^{-1} g^{kl} \partial_l \phi \partial_k f g_{\alpha\beta}, \tag{9}$$

$$\nabla_{i\alpha}^2 f = 0. \tag{10}$$

Hence

$$\begin{aligned}
\left|\widetilde{\nabla}^2 f - \frac{\widetilde{g}}{2t}\right|^2 &= \left|\nabla^2 f - \frac{g}{2t}\right|^2 + \left|\widetilde{\nabla}_{\alpha\beta}^2 f - \frac{g_{\alpha\beta}}{2t}\right|^2 \\
&= \left|\nabla^2 f - \frac{g}{2t}\right|^2 + \left|\frac{g^{kl} \partial_l \phi \partial_k f g_{\alpha\beta}}{q} + \frac{g_{\alpha\beta}}{2t}\right|^2 \\
&= \left|\nabla^2 f - \frac{g}{2t}\right|^2 + \left|\left(\frac{\nabla\phi \cdot \nabla f}{m-n} + \frac{1}{2t}\right) g_{\alpha\beta}\right|^2 \\
&= \left|\nabla^2 f - \frac{g}{2t}\right|^2 + \frac{1}{m-n}\left(\nabla\phi \cdot \nabla f + \frac{m-n}{2t}\right)^2.
\end{aligned} \tag{11}$$

The following result was obtained in a private discussion by Bing-Long Chen and the second author in January 2006.

Theorem 2.1 *([9]) The Laplace-Beltrami operator on* $(\widetilde{M}, \widetilde{g})$ *is given by*

$$\Delta_{\widetilde{M}} = L + e^{-\frac{2\phi}{m-n}} \Delta_N.$$

Proof. The proof can be given by a direct calculation. $\square$

A new proof of Theorem 1.1. To avoid technical issue, we only consider the case of compact manifolds. Let $u = \frac{e^{-f}}{(4\pi t)^{m/2}} : M \to [0, \infty)$ be a positive solution to the heat equation $\partial_t u = Lu$. Then it satisfies the following heat equation on $(\widetilde{M}, \widetilde{g})$

$$\partial_t u = \Delta_{\widetilde{M}} u.$$

Since f depends only on the variable in the M-direction, we have $\widetilde{\nabla} f = \nabla f$. Therefore the W-entropy functional $W_m(u,t)$ defined by (2) coincides with the W-entropy functional $\widetilde{W}_m(u,t)$ defined on $(\widetilde{M}, \widetilde{g})$ as follows

$$\widetilde{W}_m(u,t) = \int_{\widetilde{M}} \left[t|\widetilde{\nabla} f|^2 + f - m\right] \frac{e^{-f}}{(4\pi t)^{m/2}} dvol_{\widetilde{M}}. \tag{12}$$

Applying the W-entropy formula for the heat equation $\partial_t u = \Delta u$ on compact Riemannian manifolds with fixed metric due to Ni [28, 29] to $(\widetilde{M}, \widetilde{g})$, we have

$$\frac{d\widetilde{W}_m(u,t)}{dt} = -2\int_{\widetilde{M}} t\left(\left|\widetilde{\nabla}^2 f - \frac{\widetilde{g}}{2t}\right|^2 + \widetilde{Ric}(\widetilde{\nabla}\log u, \widetilde{\nabla}\log u)\right) u d\mu dv_N. \tag{13}$$

By (11), we have

$$\left|\widetilde{\nabla}^2 f - \frac{\widetilde{g}}{2t}\right|^2 = \left|\nabla^2 f - \frac{g}{2t}\right|^2 + \frac{2}{m-n}\left(\nabla\phi\cdot\nabla f + \frac{m-n}{2t}\right)^2. \tag{14}$$

On the other hand, by [6, 21, 17], we have

$$\widetilde{Ric}(\widetilde{\nabla}\log u, \widetilde{\nabla}\log u) = Ric_{m,n}(L)(\nabla\log u, \nabla\log u). \tag{15}$$

From (13), (14) and (15), we obtain (3). This finishes the new proof of Theorem 1.1. $\square$

Remark 2.2 One of the advantages of the above proof is that: when $m \in \mathbb{N}$ and $m > n$, the quantity $\frac{1}{m-n}\left(\nabla\phi\cdot\nabla f + \frac{m-n}{2t}\right)^2$ appeared in the W-entropy formula in Theorem 1.1 has a natural geometric interpretation. It corresponds to the vertical component of the quantity $\left|\widetilde{\nabla}^2 f - \frac{\widetilde{g}}{2t}\right|^2$ on the warped product manifold $\mathcal{M} = M \times N$ equipped with the metric

$$\widetilde{g} = g \bigoplus e^{-\frac{2\phi}{m-n}} g_N.$$

3 Dissipation formula of the Boltzmann-Shannon entropy

Let $(M, g(t), f(t))$ be as in Theorem 1.2. Following [2, 21, 17], we introduce the Bakry-Emery Ricci curvature associated with L as

$$Ric(L) = Ric + \nabla^2 f.$$

The purpose of this section is to prove the following dissipation formula for the Boltzmann-Shannnon entropy associated with the Witten Laplacian on manifolds with time dependent metrics and potentials.

Theorem 3.1 *Let u be a positive solution to the heat equation $\partial_t u = Lu$. Let*

$$H(u,t) = -\int_M u \log u d\mu$$

be the Boltzmann-Shannon entropy associated with the Witten Laplacian L. Then

$$\frac{\partial^2}{\partial t^2} H(u,t) = -2\int_M \left[|\nabla^2 \log u|^2 + \left(\frac{1}{2}\frac{\partial g}{\partial t} + Ric(L)\right)(\nabla \log u, \nabla \log u)\right] u d\mu. \tag{16}$$

Proof. By direct calculation, we have

$$\frac{\partial}{\partial t} H(u,t) = -\int_M \partial_t u(\log u + 1) d\mu = -\int_M Lu(\log u + 1) d\mu.$$

Integrating by parts yields

$$\frac{\partial}{\partial t} H(u,t) = \int_M |\nabla \log u|^2_{g(t)} u d\mu,$$

which further implies that, as $\partial_t(d\mu) = 0$, we have

$$\begin{aligned}
\frac{\partial^2}{\partial t^2} H(u,t) &= \int_M \frac{\partial}{\partial t}(|\nabla \log u|^2_{g(t)} u) d\mu \\
&= \int_M \left[\frac{\partial}{\partial t} g^{ij} \nabla_i \log u \nabla_j \log u\right] u d\mu + \int_M \frac{\partial}{\partial t}\left[\frac{|\nabla u|^2}{u}\right]_{\mathrm{g(t)\ fixed}} d\mu \\
&= \int_M \left[-\frac{\partial}{\partial t} g_{ij} \nabla_i \log u \nabla_j \log u\right] u d\mu + \int_M \frac{\partial}{\partial t}\left[\frac{|\nabla u|^2}{u}\right]_{\mathrm{g(t)\ fixed}} d\mu \\
&= \int_M \left(-\frac{\partial g}{\partial t}(\nabla \log u, \nabla u) + \frac{\partial}{\partial t}\left[\frac{|\nabla u|^2}{u}\right]_{\mathrm{g(t)\ fixed}}\right) d\mu, \qquad (17)
\end{aligned}$$

where $[\cdot]_{\mathrm{g(t)\ fixed}}$ means that the quantity $|\nabla u|^2$ in $[\cdot]$ is defined under a fixed metric $g(t)$. and we have used the facts $|\nabla \log u|^2 = g^{ij}\nabla_i \log u \nabla_j \log u$ as well as $\partial_t g^{ij} = -\partial_t g_{ij}$.

By the entropy dissipation formula in [2, 20], we have

$$\int_M \frac{\partial}{\partial t}\left[\frac{|\nabla u|^2}{u}\right]_{\mathrm{g(t)\ fixed}} d\mu = -2\int_M \left[|\nabla^2 \log u|^2 + Ric(L)(\nabla \log u, \nabla \log u)\right] u d\mu. \tag{18}$$

Combining (17) and (18), we finish the proof of Theorem 3.1. □

As an easy consequence of Theorem 1.2, we have

Corollary 3.2 *Let $(M, g(t))$ be a closed manifold with a potential $f(t)$. Suppose that $(g(t), f(t))$ satisfies Perelman's super Ricci flow and the conjugate equation, i.e.,*

$$\begin{aligned}
\frac{\partial g}{\partial t} &\geq -2Ric(L), \\
\frac{\partial f}{\partial t} &= \frac{1}{2}\mathrm{Tr}\left(\frac{\partial g}{\partial t}\right).
\end{aligned}$$

Let u be a positive solution to the heat equation $\partial_t u = Lu$. Then the Boltzmann-Shannon entropy

$$H(u,t) = -\int_M u \log u d\mu$$

is concave in time t, i.e.,

$$\frac{d^2}{dt^2} H(u,t) \leq 0.$$

4 Proofs of Theorem 1.2 and Theorem 1.3

Following [20], we introduce

$$W(u,t)=\frac{d}{dt}(tH(u,t)).$$

By direct calculation, we can prove the following

Proposition 4.1 *We have*

$$W(u,t)=\int_M \left[t|\nabla \log u|^2-\log u\right] ud\mu,$$

and

$$\begin{aligned}\frac{d}{dt}W(u,t) &= -2\int_M t\left[|\nabla^2\log u|^2+\left(\frac{1}{2}\frac{\partial g}{\partial t}+Ric(L)\right)(\nabla\log u,\nabla\log u)\right]ud\mu\\ &\qquad +2\int_M |\nabla\log u|^2 ud\mu. \qquad (19)\end{aligned}$$

Remark 4.2 From (19), we can derive that, if

$$\frac{1}{2}\frac{\partial g}{\partial t}+Ric(L)-\frac{1}{t}\geq 0.$$

Then

$$\frac{dW(u,t)}{dt}\leq 0.$$

Let

$$H_m(u,t)=-\int_M u\log u d\mu-\frac{m}{2}(1+\log(4\pi t)).$$

Following [30, 28, 18, 20], we define $W_m(u,t)$ by the Boltzmann entropy formula

$$W_m(u,t)=\frac{d}{dt}(tH_m(u)). \qquad (20)$$

We can verify that $W_m(u,t)$ coincides with the expression given in Theorem 1.2, i.e.,

$$W_m(u,t)=\int_M \left[t|\nabla\log u|^2-\log u\right]ud\mu-\frac{m}{2}(2+\log(4\pi t)).$$

Proof of Theorem 1.2. By (20) and (16) in Theorem 3.1, we have

$$\begin{aligned}\frac{dW_m(u)}{dt} &= -2\int_M t\left[|\nabla^2\log u|^2+\left(\frac{1}{2}\frac{\partial g}{\partial t}+Ric(L)\right)(\nabla\log u,\nabla\log u)\right]ud\mu\\ &\qquad +2\int_M |\nabla\log u|^2 ud\mu-\frac{m}{2t}. \qquad (21)\end{aligned}$$

Note that

$$2t|\nabla^2\log u|^2+\frac{m}{2t}=2t\left|\nabla^2\log u+\frac{g}{2t}\right|^2+\frac{m-n}{2t}-2\Delta\log u.$$

Hence

$$\begin{aligned}\frac{d}{dt}W_m(u,t) &= -\frac{m-n}{2t}-2t\int_M\left|\nabla^2\log u+\frac{g}{2t}\right|^2ud\mu+2\int_M|\nabla\log u|^2ud\mu+2\int_M\Delta\log u ud\mu\\ &\quad -2\int_M t\left(\frac{1}{2}\frac{\partial g}{\partial t}+Ric(L)\right)(\nabla\log u,\nabla\log u)ud\mu.\end{aligned}$$

Integrating by part yields

$$\begin{aligned}\int_M\Delta\log u ud\mu &= \int_M(L\log u+\nabla f\cdot\nabla\log u)ud\mu\\ &= -\int_M|\nabla\log u|^2ud\mu+\int_M\nabla f\cdot\nabla\log u ud\mu,\end{aligned}$$

whence

$$\begin{aligned}\frac{d}{dt}W_m(u,t) &= -\frac{m-n}{2t}-2t\int_M\left|\nabla^2\log u+\frac{g}{2t}\right|^2ud\mu+2\int_M\nabla f\cdot\nabla\log u ud\mu\\ &\quad -2\int_M t\left(\frac{1}{2}\frac{\partial g}{\partial t}+Ric(L)\right)(\nabla\log u,\nabla\log u)ud\mu.\end{aligned}$$

Note that

$$\begin{aligned}&\frac{m-n}{2t}+2tRic(L)(\nabla\log u,\nabla\log u)-2\nabla f\cdot\nabla\log u\\ =\ & 2tRic_{m,n}(L)(\nabla\log u,\nabla\log u)+\frac{2t}{m-n}\left(\nabla f\cdot\nabla\log u-\frac{m-n}{2t}\right)^2.\end{aligned}$$

Hence

$$\begin{aligned}\frac{d}{dt}W_m(u,t) &= -2t\int_M\left|\nabla^2\log u+\frac{g}{2t}\right|^2ud\mu-\frac{2t}{m-n}\int_M\left(\nabla f\cdot\nabla\log u-\frac{m-n}{2t}\right)^2ud\mu\\ &\quad -2\int_M t\left(\frac{1}{2}\frac{\partial g}{\partial t}+Ric_{m,n}(L)\right)(\nabla\log u,\nabla\log u)ud\mu.\end{aligned}$$

This proves the W-entropy formula in Theorem 1.2. The monotonicity result follows. The proof of Theorem 1.2 is completed. □

Proof of Theorem 1.3. The proof is similar to Perelman's monotonicity theorem for the μ-invariant for Ricci flow [30]. See also [18]. By definition, we have

$$\mu(t)=\inf_u\left\{\int_M\left[4t|\nabla u|^2-u^2\log u^2-mu^2\right](4\pi t)^{-\frac{m}{2}}d\mu\right\},\tag{22}$$

where inf is taken among all the u such that

$$\int_M(4\pi t)^{-\frac{m}{2}}u^2d\mu=1.$$

Indeed, $\mu(t)$ is the optimal constant in the following logarithmic Sobolev inequality: for all u satisfying the above condition,

$$\int_M u^2\log u^2(4\pi t)^{-\frac{m}{2}}d\mu\le-\mu(t)-m+4\int_M t|\nabla u|^2(4\pi t)^{-\frac{m}{2}}d\mu.$$

By a similar argument as used in Perelman [30] and [7, 15, 25], we can prove that the minimization problem (22) has a non-negative minimizer $u\in H^1(M,\mu)$, which satisfies the Euler-Lagrange equation

$$-4tLu-2u\log u-mu=\mu(t)u.$$

By the regularity theory of elliptic PDEs, we have $u \in C^{1,\alpha}(M)$. By an argument due to Rothaus [31], we can further prove that u is strictly positive and smooth. Hence $v = -2\log u$ is also smooth. Moreover, as a consequence of Theorem 1.2, we can derive that $\mu(t)$ is a decreasing function in t on $[0, T]$ provided that $\{g(t), f(t), t \in [0, T]\}$ satisfies the m-dimensional Perelman's super Ricci flow (3) and the conjugate equation (4). The proof of Theorem 1.3 is completed. □

Remark 4.3 Let $m \in \mathbb{N}$ and $m > n$. Let (N, g_N) be a compact Riemniann manifold of dimension $q = m - n$. Let $\mathcal{M} = M \times N$ be the product manifold equipped with the time dependent warped product metric

$$\widetilde{g}(t) = g(t) \bigoplus e^{-\frac{2f(t)}{m-n}} g_N.$$

Similarly to Remark 2.2, the quantity $\frac{1}{m-n}\left(\nabla f \cdot \nabla \log u - \frac{m-n}{2t}\right)^2$ appeared in the W-entropy formula in Theorem 1.2 has a natural geometric interpretation. It corresponds to the vertical component of the quantity $\left|\widetilde{\nabla}^2 \log u + \frac{\widetilde{g}}{2t}\right|^2$ on $(\mathcal{M}, \widetilde{g}(t))$.

5 The W-entropy for the Ricci flow on warped product manifolds

Let $m \in \mathbb{N}$ and $m \geq n$. Let $\mathbb{T}^q$ be the q-dimensional torus with a fixed flat metric given in local coordinates by $\sum_{i=1}^q dx_i^2$, where $q = m - n$. Let $\widetilde{M} = M \times \mathbb{T}^q$ be equipped with a time dependent warped product metric

$$\widetilde{g}(t) = \sum_{i,j=1}^n g_{ij}(t) dx^i dx^j + u(t)^{\frac{2}{q}} \sum_{\alpha=1}^q dx_\alpha^2.$$

In [22], Lott studied the Ricci flow $\widetilde{g}(t)$ on the warped product manifold $\widetilde{M} = M \times \mathbb{T}^q$, which consists of a modified Ricci flow for the Riemannian metric $g(t)$ and a forward heat equation for a potential function $\psi(t) = -\log u(t)$ on the manifold M. In this section we use the Perelman's W-entropy formula for the Ricci flow $\widetilde{g}(t)$ on the warped product manifold $\widetilde{M}$ to derive the W-entropy formula for the backward heat equation associated with the Witten Laplacian $L = \Delta_{g(t)} - \nabla_{g(t)}\psi(t) \cdot \nabla_{g(t)}$ on the compact manifold M equipped with Lott's modified Ricci flow $g(t)$ and the time dependent potential $\psi(t)$.

We first recall Lott's Ricci flow on $\widetilde{M} = M \times \mathbb{T}^q$. Let $u = e^{-\psi}$. Let Ric_q be the Ricci curvature on $(\widetilde{M}, \widetilde{g})$, and Ric the Ricci curvature on (M, g). By calculation on warped product manifolds, see [6, 21, 22], we have

$$Ric_q = Ric_\psi^q + \frac{1}{q}(\Delta\psi - |\nabla\psi|^2) u^{\frac{2}{q}} \sum_{\alpha=1}^q dx_\alpha^2, \tag{23}$$

where Ric_ψ^q is the m-dimensional Bakry-Emery Ricci curvature on (M, g) with respect to the potential function ψ, i.e.,

$$Ric_\psi^q := Ric + Hess\psi - \frac{1}{q}\nabla\psi \otimes \nabla\psi.$$

See [2, 18, 19, 20]. By (23), the scalar curvature on $(\widetilde{M}, \widetilde{g})$, denoted by R_q, is given by

$$R_q = R + 2\Delta\psi - \left(1 + \frac{1}{q}\right)|\nabla\psi|^2.$$

By definition, the Ricci flow on $\widetilde{M}$ is defined by

$$\partial_t \widetilde{g} = -2Ric_q. \tag{24}$$

According to Lott [22], the Ricci flow equation (24) is equivalent to the following equations

$$\begin{aligned} \partial_t g &= -2Ric_\psi^q, && (25) \\ \partial_t \psi &= \Delta\psi - |\nabla\psi|^2. && (26) \end{aligned}$$

Note that, the first equation (25) is indeed a modified Ricci flow equation for the metric $g(t)$ on M, and the second one (26) is a forward heat equation for the potential function $\psi(t)$ on $(M, g(t))$. The systems (25) and (26) are different from Perelman's Ricci flow and the conjugate heat equation introduced in [30], i.e.,

$$\begin{aligned} \frac{\partial g}{\partial t} &= -2(Ric + \nabla^2 f), \\ \frac{\partial f}{\partial t} &= -\Delta f - R, \end{aligned}$$

and are also different from the following m-dimensional Perelman's Ricci flow and the conjugate heat equation

$$\begin{aligned} \frac{\partial g}{\partial t} &= -2(Ric + \nabla^2 f - \frac{\nabla f \otimes \nabla f}{m-n}), \\ \frac{\partial f}{\partial t} &= -\Delta f + \frac{|\nabla f|^2}{m-n} - R. \end{aligned}$$

Let ϕ be a positive solution to the conjugate heat equation on $(\widetilde{M}, \widetilde{g})$

$$\partial_t \phi = -\Delta_{\widetilde{M}}\phi + R_q \phi. \tag{27}$$

Let $\tau \in [0, T]$ be such that

$$\partial_t \tau = -1.$$

Write

$$\phi = (4\pi\tau)^{-\frac{n+q}{2}} e^{-\eta}.$$

Then

$$\partial_t \eta = -\Delta_{\widetilde{M}}\eta + |\nabla\eta|^2 - R_q + \frac{n+q}{2\tau},$$

Following Perelman [30], the W-entropy for the Ricci flow $\widetilde{g}(t)$ on the warped product manifold $\widetilde{M}$ is defined as follows

$$W(\widetilde{g}, \eta, \tau) = \int_{\widetilde{M}} [\tau(|\widetilde{\nabla}\eta|^2_{\widetilde{M}} + R_q) + \eta - (n+q)]\phi dvol_{\widetilde{M}},$$

where $dvol_{\widetilde{M}} = u dvol_M dvol_{\mathbb{T}^q}$ is volume form on $(\widetilde{M}, \widetilde{g})$.

Applying Perelman's W-entropy formula for the Ricci flow [30] to $(\widetilde{M}, \widetilde{g})$, we have

$$\frac{d}{d\tau} W(\widetilde{g}, \eta, \tau) = -2\tau \int_{\widetilde{M}} \left| Ric_q + \widetilde{Hess}\eta - \frac{\widetilde{g}}{2\tau} \right|^2_{\widetilde{M}} \phi dvol_{\widetilde{M}}. \tag{28}$$

By Theorem 2.1, the Laplace-Beltrami on $(\widetilde{M}, \widetilde{g})$ is given by

$$\Delta_{\widetilde{M}} = L + u^{-\frac{2}{q}} \Delta_{\mathbb{T}^q},$$

where

$$L = \Delta - \nabla\psi \cdot \nabla.$$

Here Δ and ∇ are respectively the Laplace-Beltrami operator and the gradient operator on (M, g). In the case ϕ is a function depending only on the variable of horizontal direction, the conjugate heat equation (27) turns out to be the following backward heat equation associated with the Witten Laplacian on $(M, g(t))$

$$\partial_t \phi = -L\phi + R_q \phi. \tag{29}$$

In this case, η is a function depending only on variable in M. Thus

$$\begin{aligned} W(\widetilde{g}, \eta, \tau) &= \int_{M\times\mathbb{T}^q} [\tau(|\nabla\eta|^2 + R_q) + \eta - (n+q)]\phi u dvol_M dvol_{\mathbb{T}^q} \\ &= \int_M \left[\tau(|\nabla\eta|^2 + R + 2\Delta\psi - \left(1 + \frac{1}{q}\right)|\nabla\psi|^2) + \eta - (n+q)\right]\phi d\mu. \end{aligned}$$

Here $d\mu = udvol_M$, and we assume $vol(\mathbb{T}^q) = 1$. Note that, for any vector field v on $\widetilde{M}$, by (8), (9) and (10), we have

$$\widetilde{Hess} v = Hess v - \frac{1}{q}u^{\frac{2}{q}}\langle\nabla\psi, \nabla v\rangle \sum_{\alpha=1}^{q} dx_\alpha^2. \tag{30}$$

Substituting (23) and (30) into (28), we have

$$\begin{aligned} \frac{d}{d\tau}W(\widetilde{g}, \eta, \tau) &= -2\tau \int_M \left| Ric_\psi^q + Hess\eta - \frac{g}{2\tau} + \frac{1}{q}(\Delta\psi - |\nabla\psi|^2 - \langle\nabla\psi, \nabla\eta\rangle - \frac{q}{2\tau})u^{\frac{2}{q}} \sum_{\alpha=1}^{q} dx_\alpha^2 \right|_{\widetilde{M}}^2 \phi d\mu \\ &= -2\tau \int_M \left[\left| Ric_\psi^q + Hess\eta - \frac{g}{2\tau} \right|^2 + \frac{1}{q}(\Delta\psi - |\nabla\psi|^2 - \langle\nabla\psi, \nabla\eta\rangle - \frac{q}{2\tau})^2 \right] \phi d\mu. \end{aligned}$$

Thus we have proved the following W-entropy formula for the backward heat equation associated with the Witten Laplacian on compact manifolds equipped with Lott's modified Ricci flow and time dependent potentials.

Theorem 5.1 *Let $(M, g(t), \psi(t))$ be a compact manifold with a family of Remannian metrics $g(t)$ and potentials $\psi(t)$ which satisfy*

$$\begin{aligned} \partial_t g &= -2(Ric + Hess\psi - \frac{1}{q}\nabla\psi \otimes \nabla\psi), \\ \partial_t \psi &= \Delta\psi - |\nabla\psi|^2. \end{aligned}$$

Let $d\mu = e^{-\psi} dvol_M$, and $L = \Delta - \nabla\psi \cdot \nabla$. Let ϕ be a positive solution to the backward heat equation of the Witten Laplacian on M, i.e.,

$$\partial_t \phi = -L\phi + R_q \phi.$$

where $R_q = R + 2\Delta\psi - (1 + \frac{1}{q})|\nabla\psi|^2$. Let

$$H_{n+q}(\phi, \tau) = -\int_M \phi \log\phi d\mu - \frac{n+q}{2}(1 + \log(4\pi\tau)),$$

and define the W-entropy $W_q(g, \psi, \eta, \tau)$ by

$$W_q(g, \psi, \eta, \tau) = \frac{d}{d\tau}(\tau H_{n+q}(\phi, \tau)).$$

Then

$$W_q(g,\psi,\eta,\tau)=\int_M[\tau(|\nabla\eta|^2+R_q)+\eta-(n+q)]\phi d\mu.$$

where $\phi=(4\pi\tau)^{-\frac{n+q}{2}}e^{-\eta}$. *Moreover*

$$\frac{d}{d\tau}W_q(g,\psi,\eta,\tau)=-2\tau\int_M\left[\left|Ric_\psi^q+Hess\eta-\frac{g}{2\tau}\right|^2+\frac{1}{q}\left(\Delta\psi-|\nabla\psi|^2-\langle\nabla\psi,\nabla\eta\rangle-\frac{q}{2\tau}\right)^2\right]\phi d\mu.$$

In particular, $W_q(g,\psi,\eta,\tau)$ *is decreasing in the backward time* τ*, and the monotonicity is strict unless that*

$$\begin{aligned}Ric_\psi^q+Hess\eta &= \frac{g}{2\tau},\\ \Delta\psi-|\nabla\psi|^2 &= \langle\nabla\psi,\nabla\eta\rangle-\frac{q}{2\tau}.\end{aligned}$$

As an application of the W-entropy formula in Theorem 1.3, we have the following result, which is the restatement of Perelman's monotonicity theorem for the μ-invariant for Ricci flow [30] on warped product manifolds. See also [7, 10, 15, 25].

Theorem 5.2 *Let* $(M,g(t),\psi(t))$ *be a compact manifold with a family of Remannian metrics* $g(t)$ *and potentials* $\psi(t)$ *which satisfy*

$$\begin{aligned}\partial_t g &= -2(Ric+Hess\psi-\frac{1}{q}\nabla\psi\otimes\nabla\psi),\\ \partial_t\psi &= \Delta\psi-|\nabla\psi|^2.\end{aligned}$$

Then there exists a positive and smooth function $u=e^{-\frac{\eta}{2}}$ *such that* η *achieves the optimal logarithmic Sobolev constant* $\mu(\tau)$ *defined by*

$$\mu(\tau):=\inf\left\{W_q(g,\psi,\eta,\tau):\ \int_M\frac{e^{-\eta}}{(4\pi\tau)^{\frac{n+q}{2}}}d\mu=1\right\}.$$

where

$$W_q(g,\psi,\eta,\tau)=\int_M[\tau(|\nabla\eta|^2+R_q)+\eta-(n+q)]\phi d\mu,$$

Indeed, $u=e^{-\frac{\eta}{2}}$ *is a solution to the nonliniear PDE*

$$-4\tau Lu+\tau R_q u-2u\log u-(n+q)u=\mu(\tau)u.$$

Moreover, $\mu(\tau)$ *is decreasing in* τ *on* $[0,T]$.

Acknowledgement. The second author would like to thank Bing-Long Chen for useful discussion in January 2006.

References

[1] B. Andrews, L. Ni, Eigenvalue comparison on Bakry-Emery manifolds, to appear in Comm. Partial Differential Equations. arxiv1111.4967v1

[2] D. Bakry, M. Emery, Diffusion hypercontractives, in: Sém. in Probab. XIX, in: Lecture Notes in Math., vol. 1123, Springer-Verlag, Berlin/New York, 1985, 177-206.

[3] D. Bakry, M. Ledoux, A logarithmic Sobolev form of the Li-Yau parabolic inequality, Rev. Mat. Iberoamericana, 22(2) (2006), 683-702.

[4] D. Bakry, Z.-M. Qian, Harnack inequalities on a manifold with positive or negative Ricci curvature, Rev. Mat. Iberoamericana 15 (1999) 143-179.

[5] D. Bakry, Z.-M. Qian, Some new results on eigenvectors via dimension, diameter, and Ricci curvature, Adv. in Math. 155 (2000), 98-153.

[6] A. Besse, Einstein Manifolds, Springer, Berlin, 1987.

[7] H.-D. Cao, X.-P. Zhu, A complete proof of the Poincaré and geometrization conjecturesapplication of the Hamilton-Perelman theory of the Ricci flow. Asian J. Math. 10(2), 165-492 (2006). Erratum to: A complete proof of the Poincar and geometrization conjectures-application of the Hamilton-Perelman theory of the Ricci flow [Asian J. Math. 10(2), 165492] (2006); Asian J. Math. 10(4), 663 (2006)

[8] S.-C. Chang, T.-J. Kuo, S.-H. Lai, Li-Yau gradient estimate and entropy formulae for the CR heat equation in a closed pseudohermitian 3-manifold, J. Diff. Geom. 89, No. 2 (2011), 185-216.

[9] B.-L. Chen, X.-D. Li, Private discussion in January 2006.

[10] B. Chow, P. Lu, L. Ni, Hamilton's Ricci Flow, Lectures in Comtemporary Maths., Sciences Press, Beijing, Amer. Math. Soc. 2006.

[11] K. Ecker, A formula relating entropy monotonicity to Harnack inequalities, Comm. Anal. Geom. 15 (2007), no. 5, 1025–1061.

[12] F. Fang, X.-D. Li, Z.-L. Zhang, Two generalizations of Cheeger-Gromoll splitting theorem via Bakry Emery Ricci curvature. Ann. Inst. Fourier (Grenoble) 59(2) (2009), 563-573.

[13] A. Futaki, H.-Z. Li, X.-D. Li, On the first eigenvalue of the Witten Laplacian and the diameter of compact shrinking solitons, Ann. Glob. Anal. Geom. DOI 10.1007/s10455-012-9358-5.

[14] R.S. Hamilton, A matrix Harnack estimate for the heat equation, Comm. Anal. Geom. 1 (1993), no. 1, 113-126.

[15] B. Kleiner, J. Lott, Notes on Perelman's papers, Geom. Topol. 12 (2008), no. 5, 2587-2855.

[16] P. Li, S.-T. Yau, On the parabolic kernel of the Schrödinger operator, Acta. Math. 156 (1986), 153-201.

[17] X.-D. Li, Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds, J. Math. Pures Appl. 84 (2005), 1295-1361.

[18] X.-D. Li, Perelman's entropy formula for the Witten Laplacian on Riemannian manifolds via Bakry-Emery Ricci curvature, Math. Ann. 353 (2012), 403-437.

[19] X.-D. Li, Perelman's W-entropy for the Fokker-Planck equation over complete Riemannian manifolds, Bull. Sci. math. 135 (2011) 871-882.

[20] X.-D. Li, Harnack inequality, gradient estimates and W-entropy formula on complete Riemannian manifolds, arxiv:1303.1242v1.

[21] J. Lott, Some geometric properties of the Bakry-Emery Ricci tensor, Comment. Math. Helv. 78 (2003) 865-883.

[22] J. Lott, Optimal Transport and Perelman's Reduced Volume, Calc. Var. and Partial Differential Equations 36 (2009), 49-84.

[23] P. Lu, L. Ni, J.-J. Vazquez, C. Villani, Local Aronson-Benilan estimates and entropy formulae for porous medium and fast diffusion equations on manifolds, J. Math. Pures Appl., Vol 91 (2009), no. 1, 1-19.

[24] B. Kotschwar, L. Ni, Gradient estimate for p-harmonic functions, $1/H$ flow and an entropy formula, Ann. Sci. Ec. Norm. Sup., Vol 42 (2009), no. 1, 1–36.

[25] J. W. Morgan, G. Tian, Ricci flow and the Poincar conjecture. Clay Mathematics Monographs, 3. American Mathematical Society, Providence, Clay Mathematics Institute, Cambridge (2007)

[26] F. Otto, The geometry of dissipative evolution equations: the porous medium equation, Commun. in Parial Differential Equations, 26 (1 and 2) (2001), 101-174.

[27] F. Otto and C. Villani, Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality, J. Funct. Anal. 173 (2000) 361-400.

[28] L. Ni, The entropy formula for linear equation, J. Geom. Anal. 14 (1) (2004), 87-100.

[29] L. Ni, Addenda to "The entropy formula for linear equation", J. Geom. Anal. 14 (2) (2004), 329-334.

[30] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, http://arXiv.org/abs/maths0211159.

[31] O.S. Rothaus, Logarithmic Sobolev inequalities and the spectrum of Schrodinger opreators, J. Funct. Anal., 42:1 (1981), 110-120.

[32] K.-T. Sturm, Convex functionals of probability measures and nonlinear diffusions on manifolds, J. Math. Pures Appl. (9) 84 (2005), no. 2, 149-168.

[33] K.-T. Sturm and M.-K. von Renesse, Transport inequalities, gradient estimates, entropy, and Ricci curvature Comm. Pure Appl. Math., 58(7) (2005), 923–940.

[34] C. Villani, Topics in Mass Transportation, Grad. Stud. Math., Amer. Math. Soc., Providence, RI, 2003.

[35] C. Villani, Optimal Transport, Old and New, Springer, 2008.

[36] G. Wei, W. Wylie, Comparison geometry for the Bakry-Emery Ricci tensor, J. Diff. Geom. 83 (2009), 377-405.

Songzi Li

School of Mathematical Science, Fudan University, 220, Handan Road,
Shanghai, 200432, China

Xiang-Dong Li

Academy of Mathematics and Systems Science, Chinese Academy of Sciences,
55, Zhongguancun East Road, Beijing, 100190, China, E-mail: xdli@amt.ac.cn